# **Elektrolyt**

## agilan® complementary feed

To support the supply with minerals and active components, water-soluble

For poultry, pigs, calves, cattle and horses

### Composition

Glucose

Monoammonium phosphate

Sodium chloride

Sodium bicarbonate

Potassium chloride

### **Analytical constituents**

Crude protein	6.5 %
Crude fat	0.1 %
Crude fibre	0.0 %
Crude ash	33.0 %
Ash insoluble in HCl	3.0 %
Calcium	3.0 %
Phosphorus	2.5 %
Sodium	4.5 %
Lysine	0.0 %
Methionine	0.0 %



Prevention of nutrition-related disorders. Deficiencies of minerals and vitamins which lead to metabolic disturbances and may cause stress situations.

### Content of additives per kg

#### Vitamins.

<u>vicaninis.</u>	
Vitamin D <sub>3</sub> (3a671)	3,000 I.U.
Vitamin E (3a700)	1,200 mg
Vitamin B1 (3a821)	45 mg
Vitamin B2 (3a825i)	75 mg
Vitamin B6 (3a831)	45 mg
Vitamin B12 / Cyanocobalamin	150 µg
Vitamin K <sub>3</sub> (3a711)	300 mg
Vitamin C (3a300)	3,200 mg
Niacinamid (3a315)	450 mg
Calcium-D- pantothenate (3a841)	150 mg
Folic acid (3a316)	15 mg
Biotin (3a880)	3,000 µg

















# **Elektrolyt**

### Trace elements:

Zinc (3b605)	(in form of zinc sulphate monohydrate)	800 mg
Manganese (3b503)	(in form of manganous sulphate, monohydrate)	600 mg
Iron (3b103)	(in form of iron (II) sulphate, monohydrate)	400 mg
Copper (3b405)	(in form of copper (II) sulphate, pentahydrate)	80 mg
Iodine (3b202)	(in form of calcium iodate, anhydrous)	130 mg
Selenium (3b801)	(in form of sodium selenite)	3 mg

### Feeding recommendation

20 g / litre drinking water

### **Feeding instruction**

Due to the increased level of trace elements compared to complete feed, this complementary feed may only be used up to 3.8 % of the daily ration fed to poultry, pigs, calves, cattle and horses.

Simultaneous use with Vitamin D2 is not allowed.

#### Storage

Shelf life guarantee is only valid for unopened packages, stored at a cool and dry place.

Date of revision of the text: August 2022

