

# Dystan®-Powder

agilan® complementary feed  
with spruce needles, wormwood, balm, chamomile and  
pure apple pectin

For all animal species

## Composition

Dextrose  
Spruce needles (11 %)  
Apple pectin (10 %)  
Sodium chloride  
Powdered milk  
Psyllium husks ground  
Sodium bicarbonate  
Potassium chloride

## Analytical constituents

|               |        |
|---------------|--------|
| Crude protein | 11.7 % |
| Crude fat     | 2.9 %  |
| Crude fibre   | 10.1 % |
| Crude ash     | 20.5 % |
| Sodium        | 4.0 %  |
| Lysine        | 0.2 %  |
| Methionine    | 0.0 %  |

## Content of additives per kg

### Trace elements:

|   |          |
|---|----------|
| Copper (3b405) (as Cu-II-sulphate, pentahydrate)            | 750 mg   |
| Zinc (3b605) (as Zn sulphate, monohydrate)                  | 4,000 mg |
| Manganese (3b503) (as manganous-(II)-sulphate, monohydrate) | 3,000 mg |

### Flavouring compounds:

|   |          |
|---|----------|
| Melissa officinalis 70/524/EEC; CoE 280 (Extract) | 4,000 mg |
| Matricaria recutita 70/524/EEC (Extract)          | 5,000 mg |
| Artemisia absinthium L. 70/524/EEC (Tincture)     | 1,000 mg |

## Recommended use

With natural ingredients such as lemon balm, chamomile and wormwood for supplementary feeding as needed for digestive disorders and intestinal problems such as a change in faecal consistency, flatulence and a general reluctance to eat.

## Feeding recommendation

The present statements apply as long as the veterinarian has not advised otherwise:

- Calves:

As needed, 2-3 x a day dissolve 50 g Dystan® Powder in 1 - 2 litres of warm water (40 °C) and add to feed (or 0.1 – 1.0 g/ kg BW [body weight]).



# Dystan<sup>®</sup>-Powder

- Cattle, horses:  
50 - 75 g per administration (or 0.1 g/ kg body weight)
- Foals:  
up to 25 g per administration (or 0.1 - 0.5 g/ kg body weight)
- Pigs (weaners, shoats):  
up to 15 g per administration (or 0.1 - 0.5 g/ kg body weight)
- Sucklings  
up to 6 g per administration (or 0.1 - 1.0 g/ kg body weight)
- Dogs, sheep:  
up to 25 g per administration (or 0.1 - 1.0 g/ kg body weight)
- Poultry:  
0.1 - 1.0 g/ kg body weight  
(do not administer through drinking water system)
- Small domestic animals:  
0.1 - 1.0 g/ kg body weight

**Do not use this product as the exclusive source of drinking water!**

## **Storage**

Store in a dry place.

Close package tightly after use so it cannot absorb any moisture. Storage life guarantee is only valid for unopened packages stored in a cool, dry place (2 - 25 °C).

Date of revision of the text: July 2022